

AD-A146 243 A SEMI-EMPIRICAL LOW LATITUDE IONOSPHERIC MODEL(U) AIR 1/1  
FORCE GEOPHYSICS LAB HANSCOM AFB MA  
D N ANDERSON ET AL. 07 SEP 84 AFGL-TR-84-0226


UNCLASSIFIED

F/G 4/1

NL



END  
DATE  
FILMED  
10 84  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

7

UNCLASSIFIED

AD-A146 243

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			12. RESTRICTIVE MARKINGS N/A													
2a. SECURITY CLASSIFICATION AUTHORITY N/A			3. DISTRIBUTION AVAILABILITY OF REPORT  Approved for public release; Distribution unlimited.													
2b. DECLASSIFICATION DOWNGRADING SCHEDULE N/A																
4. PERFORMING ORGANIZATION REPORT NUMBER AFGL-TR-84-0226			5. MONITORING ORGANIZATION REPORT NUMBER													
6a. NAME OF PERFORMING ORGANIZATION Air Force Geophysics Lab.		6b. OFFICE SYMBOL (If applicable) LIS	7a. NAME OF MONITORING ORGANIZATION													
6c. ADDRESS (City, State and ZIP Code) Hanscom AFB Massachusetts 01731			7b. ADDRESS (City, State and ZIP Code)													
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Air Force Geophysics Lab.		8b. OFFICE SYMBOL (If applicable) LIS	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER													
8c. ADDRESS (City, State and ZIP Code) Hanscom AFB, MA 01731		10. SOURCE OF FUNDING NOS. <table border="1"><tr><td>PROGRAM ELEMENT NO.</td><td>PROJECT NO.</td><td>TASK NO.</td><td>WORK UNIT NO.</td></tr><tr><td></td><td>2310</td><td>2310G9</td><td>01</td></tr></table>			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.		2310	2310G9	01				
PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.													
	2310	2310G9	01													
11. TITLE (Include Security Classification) A Semi-Empirical, Low Latitude Ionospheric Model 61102F																
12. PERSONAL AUTHOR(S) David N. Anderson, Michael Mendillo* and Bruce Herniter*																
13a. TYPE OF REPORT REPRINT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) 1984 September 7												
15. PAGE COUNT 6																
16. SUPPLEMENTARY NOTATION *Dept of Astronomy, Boston University, Boston, MA 02215 Presented at the 7th Intern'l Symposium on Equatorial Aeronomy, 22-29 March 1984, Hong Kong																
17. COSATI CODES <table border="1"><tr><td>FIELD</td><td>GROUP</td><td>SUB. GR.</td></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr></table>			FIELD	GROUP	SUB. GR.										18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Electron density, plasma transport, theoretical ionospheric model, Low latitude ionosphere.	
FIELD	GROUP	SUB. GR.														
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Currently available emperical models of the low latitude ionosphere are seriously in error, especially during high sunspot activity years. Substantial improvement in models is achieved when theoretically calculated profiles are used. However, the fastest theoretical model is prohibitively time-consuming on even the fastest computer. A unique solution to this problem is to theoretically calculate electron density profiles as a function of latitude and local time and then generate coefficients which reproduce these profiles. The coefficients themselves are then easily stored, quickly retrieved and form the basis for a fast, portable, semi-empirical computer code which will produce realistic, low-latitude F-region electron density profiles.																
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED													
22a. NAME OF RESPONSIBLE INDIVIDUAL D. ANDERSON			22b. TELEPHONE NUMBER (Include Area Code) 617-861-3220	22c. OFFICE SYMBOL LIS												


DTIC  
ELECTE  
SEP 27 1984

AFGL-84-0226

# A SEMI-EMPIRICAL, LOW-LATITUDE IONOSPHERIC MODEL

David N. Anderson  
Air Force Geophysics Laboratory  
Hanscom AFB MA 01731

Michael Mendillo and Bruce Herniter  
Department of Astronomy  
Boston University  
Boston MA 02215



Accession For	
NTIS GRA&I	X
DTIC TAB	
Unannounced Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## INTRODUCTION

Current empirical models of the low-latitude ionospheric F-region (Llewellyn and Bent, 1973; Chiu, 1975; Rawer, 1981) severely underestimate the daytime plasma density scale-height and total electron content (TEC) when compared with actual observations. As illustrated in Figure 1, the International Reference Ionosphere (IRI) described by Rawer (1981) yields daytime TEC values which are 50% lower than observed TEC values at Manila (dip, 14.5°N) in January, 1982. Some improvement occurs when the Bent topside model is incorporated with the IRI but the predicted values are still lower than observed. Substantial improvement is achieved when theoretically calculated profiles (Anderson, 1973) are used to predict TEC values. The reason is that vertical plasma transport by upward  $\bar{E} \times \bar{B}$  drift produces both topside and bottomside profiles which are much broader (thicker) than Chapman-like profiles. To illustrate this point, Figure 2 shows a comparison between a theoretically-calculated, noontime electron density profile and the predicted IRI-Bent profile.

To calculate electron density profiles whenever they are needed, however, is prohibitively time-consuming on even the fastest computer. A unique solution to this problem is to theoretically calculate electron density profiles as a function of latitude and local time and then generate coefficients which reproduce these profiles. The coefficients themselves are easily stored, quickly retrieved and

84 09 24 054

form the basis for a fast, portable, semi-empirical computer code which will produce realistic, low-latitude F-region electron density profiles.

#### DISCUSSION

Using the techniques described by Anderson (1973), electron densities as a function of altitude, latitude and local time are calculated by solving the time-dependent ion ( $O^+$ ) continuity equation numerically. The effects of production by photo-ionization, loss through charge exchange with  $N_2$  and  $O_2$  and transport by diffusion, neutral winds and vertical  $\bar{E} \times \bar{B}$  drift are included. Because  $\bar{E} \times \bar{B}$  drifts play such an important role in determining the electron density distribution and have a seasonal and solar cycle dependence, the calculations are carried out for equinox and solstice conditions for both solar cycle maximum and minimum periods. The  $\bar{E} \times \bar{B}$  drift models are patterned after Jicamarca incoherent scatter radar drift observations as reported by Fejer (1981). Other input parameters include the MSIS neutral atmosphere model (Bedin et al., 1977), loss rates determined by Torr and Torr (1979) and appropriate production rates from Torr et al. (1979). A neutral wind model is derived from a number of sources including Dickinson et al. (1981) and Sipler et al. (1983).

Once electron density profiles are obtained they are normalized to the peak electron density,  $N_{MAX}(F2)$  and the coefficients which reproduce the normalized profiles are stored. Appropriate coefficients which generate electron density profiles (200-1000 km) every half-hour local time and every  $2^\circ$  dip latitude between  $26^\circ N$  and  $26^\circ S$  dip latitude are calculated. To regenerate density profiles internally,  $N_{MAX}(F2)$  is specified using the CCIR (1978) coefficients.

Being able to quickly generate realistic electron density profiles has a number of applications. Besides providing diurnal TEC values at any specified dip latitude or slant path, this model could be used to specify the ambient

ionosphere into which chemical release experiments would be performed. This model would give the initial conditions to be used by any chemical release code to determine airglow intensities, percent plasma reduction, size of the depleted region, etc. In addition, such a model would yield flux-tube integrated Pedersen conductivity and electron content values which are important in determining and predicting low-latitude, instability growth rates. Finally, the program would be capable of supplying airglow intensities both in the vertical and at any slant viewing angle desired. It could thus determine the correlation between TEC values and say, 7774 A airglow intensity which could provide "all sky" TEC values from all-sky 7774 airglow maps.

Acknowledgments: This research was supported in part by AFGL Contract F19628-81-K0051.

# REFERENCES

- Anderson, D.N., A theoretical study of the ionospheric F-region equatorial anomaly. II. Results in the American and Asian sectors, Planet. Space Sci., 21, 421, 1973.
- Chiu, Y.T., An improved phenomenological model of ionospheric density, J. Atmos. Terr. Phys., 37, 341, 1975.
- Dickinson, R.E., E.C. Ridley, and R.G. Roble, A three-dimensional general circulation model of the thermosphere, J. Geophys. Res., 86, 1499, 1981.
- Fejer, B.G., The equatorial ionospheric electric fields - a review, J. Atmos. Terr. Phys., 43, 377, 1981.
- Hedin, A.E., C.A. Reber, G.P. Newton, N.W. Spencer, H.C. Brintar, H.G. Mayr, and W.E. Potter, A global thermospheric model based on spectrometer and incoherent scatter data, MSIS2 composition, J. Geophys. Res., 82, 2148, 1977.
- International Radio Consultative Committee (CCIR), CCIR atlas of ionospheric characteristics, Rep. 340-3 Recommendation and Reports of the CCIR, 1978, Vol. 6, International Telecommunications Union, Geneva, 1978.
- Llewellyn, S.K. and R.B. Bent, Documentation and description of the Bent Ionospheric Model, AFCL-TR-73-0657, AD772733, 1973.
- Rawer, K. in International Reference Ionosphere - IRI79. Editors J.V. Lincoln, and R.O. Cankright, World Data Center A, NOAA, Boulder, CO, 1981.
- Sipler, D.P., M.A. Biondi, and R.G. Roble, F-region neutral winds and temperatures at equatorial latitudes: Measured and predicted behavior during geomagnetically quiet conditions, Planet. Space Sci., 31, 53, 1983.
- Torr, D.G. and M.R. Torr, Chemistry of the thermosphere and ionosphere, J. Atmos. Terr. Phys., 41, 797, 1979.
- Torr, M.R., D.G. Torr, R.A. Org, and H.E. Hinteregger, Ionization frequency of thermospheric constituents as a function of solar cycle 21, Geophys. Res. Lett., 10, 771, 1979.




Fig. 1






Fig. 2

D  
B4